

HYDROPHOBIC COMPOSITIONS AND METHODS OF USE IN WATER APPLICATIONS

5 FIELD OF THE INVENTION

[0001] The present invention relates to the field of hydrophobic compositions and methods of use in water applications.

BACKGROUND OF THE INVENTION

[0002] Microbes and microbial preparations are becoming increasingly important
10 and effective in combating pests and diseases. One of the most important factors in the efficacy of microbial control of pests or diseases is the application and distribution of the microbes to the targeted pest or disease. Current techniques for the application and distribution of microbes to water-related applications are deficient and hinder the effectiveness of the application and distribution of the microbes.

15 [0003] In applications to bodies of water, lakes, rivers, streams, canals, ponds, tanks, reservoirs, and water films on the surface of plants it is difficult to evenly apply microbes over the entire surface area of the body of water or plant surface and throughout the water columns beneath the water surface. For example, it is difficult to control algae blooms, which normally occur in the top layers of water, unless there is effective
20 distribution of the microbes over the surface layers followed by adequate movement of the microbes into the upper water layers where the algae are residing.

[0004] Currently, the following methods are employed to control algae blooms and other unwanted plants and organisms. One method involves the use of chemical poisons. For example, the most commonly used algaecides include copper sulfate and
25 diuron which have detrimental environmental impact, are regulated by the EPA as well as

state agencies, require expert handling, and are considered pollutants. Their application often requires the use of a boat or other device to deliver the pesticide to the target organism as persons must spray the treatment directly onto the target organism or plant. This application is very labor intensive, time consuming, and a danger to persons
5 applying the product. These and other pesticides, herbicides, and pollutants for the control of algae or other noxious plants and organisms are constantly under heavy scrutiny by municipal, state, and federal regulators nationally.

[0005] Another current method involves the use of colorings and shadings that provide limited control of the targeted organism or plant through prevention (blocking
10 sunlight that is an essential stimulant for many algae that thrive on photosynthesis). Consequently, beneficial and ornamental water plants requiring sunlight are also deprived of needed light and their performance is also inhibited. Additionally, water clarity is sacrificed and unnatural colored water is created by the dyes, which is undesirable in many circumstances.

15 [0006] Another method involves the application of ultra-violet light in the form of a pre-filter to the area water gardening and koi ponds. Ultra-violet filtration, as it is commonly called, kills beneficial and bad bacteria alike, creating a pond more reliant on chemical fixes since natural pond balancing, nitrogen fixing, and micro-flora are destroyed. Additionally, ultra-violet filtration can not be practically applied to larger
20 bodies of water or naturally sustaining bodies of water.

[0007] Another current method involves microbial inoculation, which has generally taken two forms, that of a liquid solution of microbes and a granular blend. Neither form offers an effective distribution method. Extensive mechanical agitation is

required in order to distribute a liquid microbial inoculate throughout a body of water. Without extensive mechanical agitation, these liquid microbial formulas tend to remain concentrated at the origin of application. Additionally, liquid microbial formulas with viscosities that are greater or less than water will tend to hover or sink in a specific location. The other form of microbial inoculate currently available is granular. These granular formulas are typically similar in weight to that of beach sand. When introduced into the treatment waters, these granular formulations fall directly to the bottom. Only a very tiny percentage of the granular formulation (typically in the form of wheat germ) is left on the surface immediately following treatment. This method of granular microbial inoculation is largely ineffective, as the beneficial microbes are not effectively nor efficiently delivered to the target organism (alga, fungi, weeds, diseased fish, etc).

[0008] Each of the methods described above are deficient in their effectiveness since the composition applied to the algae, noxious weed, or target organism is concentrated at the origin of application and the poisons, antibiotics, or microbes are not evenly distributed. Further, since the application of the composition is limited to the localized area of application, only those algae or other plants or organisms residing in that localized area of the body of water will be treated sufficiently with the correct dosage. The target organisms residing in the adjacent water columns will not be treated as effectively. Thus, the current methods are inefficient for the treatment of algae, plants, fish, or fungi in water applications.

[0009] Compositions and methods for delivering microbes to water applications uniformly and efficiently over the water surface and down through the water column for the treatment of pests or diseases would offer substantial benefits over the current

techniques. Accordingly, an object of this invention is a composition for use in water applications, wherein the composition distributes microbes uniformly and efficiently over a water surface and down through the water column. Another object of the present invention is a method for applying microbes uniformly and efficiently over a water surface and down through the water column. Another object of the present invention is to deliver microbes proportionately to the target alga, weeds, or organisms by blanketing the target pest effectively with the microbe.

SUMMARY OF THE INVENTION

[0010] An object of the present invention is to provide a composition and method by which microbes can be uniformly and efficiently distributed over a water surface and released into the body of water.

[0011] One aspect of the invention is a hydrophobic composition used for the delivery of a microbial agent to water applications comprising a hydrophobic carrier and a microbial agent. Another aspect of the invention is a method of administering a microbial agent to a water application uniformly and efficiently by introducing a hydrophobic composition containing a microbe to the water application and delivering it to a target plant or organism. Another aspect of the invention is a method for improving the water quality in a water application by introducing hydrophobic compositions to the water application. Another aspect of the invention is a method for improving the water quality in a water application by introducing hydrophobic compositions containing strains of *Bacillus subtilis* to the water application.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0012] This invention relates to compositions and methods for uniformly and efficiently delivering microbial agents to water applications.

[0013] A preferred aspect of the invention is a hydrophobic composition for effectively distributing a microbial agent to a water application comprising a hydrophobic carrier and a microbial agent. A preferred hydrophobic carrier is preferably selected from a group comprising carbon minerals, leonardite, plastic, wood powder, plant materials, clay and combinations thereof. The carbon minerals are preferably selected from a group comprising coal, charcoal, graphite, carbon black, oil, straw and combinations thereof.

Most preferably, the hydrophobic carrier comprises activated carbon or charcoal specifically designed for distribution onto and into water applications. The carbon formulations also confer UV protection to the microbial agent during exposure to sunlight and act as an inoculum source for the microbial agent as it passes into the liquid medium. Carbon in various forms has been used in commercial water treatment previously, but as a filter or absorbent. The hydrophobic properties of the hydrophobic composition allow the composition to resist mixing with the water and instead form a thin film that rapidly disperses over the water surface resulting in its effective application to the water surface. Accordingly, in addition to the preferred hydrophobic carriers, one skilled in the art will appreciate that any hydrophobic or carbon composition may be employed as a hydrophobic carrier.

[0014] As used herein, a microbial agent comprises fungi, bacteria, nematodes, protozoan or combinations thereof which are effective in treating target pests or diseases. The target pests or diseases include, but are not limited to, various forms of algae and leaf diseases. Preferably the microbial agent comprises the genus *Bacillus*. Most preferably,

the microbial agent comprises various strains of *Bacillus subtilis*. In a preferred embodiment, the microbial agent is formulated onto the hydrophobic carrier to produce a material with about 4.5×10^6 cfu/g of hydrophobic composition to about 4.5×10^{10} cfu/g of hydrophobic composition.

5 [0015] As used herein, a water application includes bodies of water, lakes, rivers, canals, ponds, tanks, reservoirs and water films on the surface of plants. As would be appreciated by those of skill in the art, the present invention may be used in similar water applications where the uniform and efficient distribution of a microbial agent is desired.

10 [0016] In another aspect of the invention, the hydrophobic composition may also include a mineral fertilizer. The mineral fertilizer allows the hydrophobic composition to be applied to water films on the surface of plants, such as commercial turf. The hydrophobic composition may be added to a mineral fertilizer for application to commercial turf and similar applications. In this aspect of the invention, the hydrophobic composition may also be applied to the surface of water in the form of a water-soluble
15 bag for use as an algacide for blue-green algae (cyanobacterium), microscopic algae, filamentous algae, string algae, hair algae, and blanket weed.

20 [0017] Another aspect of the present invention is a method of delivering microbes to a water application uniformly and efficiently comprising, the step of applying a hydrophobic composition to a water application wherein the hydrophobic composition comprises a hydrophobic carrier and a microbial agent. In addition to forming a thin uniformly distributed film over the water surface, the microbial agents may be released into the body of water and move downward through the primary regions where the target pests or diseases are located during and after surface distribution. The microbial agent

then interacts with the target pests or diseases and the hydrophobic carrier serves as a reservoir for the microbial agent. In the case of bacteria, the hydrophobic carrier provides a matrix for further growth.

[0018] Another aspect of the invention is a method for improving the water
5 quality of a water application. In a preferred embodiment of the invention a hydrophobic composition is introduced to the water application. The hydrophobic composition disperses rapidly as a thin film over the surface of the water application, and downward through the primary regions of the water column. The microbial agents of the hydrophobic composition are released into the water application, thereby treating and
10 improving the water quality of the water application. In further aspects of the invention, the hydrophobic compositions treat and improve water quality by controlling algae growth, de-odorizing the water application, suppressing water eutrophication, and reducing total suspended solids, ammonia, free nitrates and phosphates in the solution of water applications. The hydrophobic compositions of this aspect of the invention are
15 delivered at rates ranging from about 0.1 g/m² to about 1000 g/m², most preferably from about 1 g/m² to about 1.5 g/m². The hydrophobic composition may also be delivered at rates ranging from about 100 cfu/m² to about 10 X 10¹⁰ cfu/m².

[0019] In another preferred embodiment of the invention, a hydrophobic composition comprising various strains of *Bacillus subtilis* is formulated with charcoal to
20 produce a hydrophobic composition with about 4.5 X 10⁶ cfu/g of hydrophobic composition to about 4.5 X 10¹⁰ cfu/g of hydrophobic composition. The hydrophobic composition may be applied to water applications to control algae growth, de-odorize the water application, suppress water eutrophication, and reduce total suspended solids,

ammonia, free nitrates and phosphates in the water applications whereby the hydrophobic composition is efficiently and uniformly dispersed over the surface of and into the water application.

5 [0020] In another aspect of the invention, the hydrophobic composition may be supplemented with ground cotton stubble.

[0021] In another aspect of the invention, *Bacillus subtilis* is used to control algae growth and improve water quality in water applications. Another embodiment of the invention is a method for treating water and controlling algae growth in water applications comprising the step of applying *Bacillus subtilis* to a water application.

10 EXAMPLE 1

[0022] Example 1 illustrates the effect of a hydrophobic composition in an algae infested water application.

[0023] Various strains of *Bacillus subtilis* were formulated with charcoal to produce a hydrophobic composition with 4.5×10^8 cfu/g of hydrophobic composition..

15 The hydrophobic composition was then applied to the surface of an algae infested pond in Riverside, California, USA at a range of rates from 0.1 -1.5 g/m². The water quality of the pond was monitored over a ten day period. On certain days during the ten day period, the algae in the pond was rated from 0 to 5, with 0 indicating the worst water quality and a high incidence of algae and 5 indicating good water quality and no
20 incidence of algae. Table 1 illustrates the results of the study.

[0024] As the rates of application of the formulation increased, the water quality increased. In addition, generally, as the rates of formulation increased, the water quality increased at faster rates.

TABLE 1

Rate of Formulation	Day 0	Day 3	Day 5	Day 7	Day 10
0g	2	2	2	2	2
0.1g	2	2	2	2	2
0.5g	2	2	3	3	3
1g	2	2	2	3	4
1.5g	2	2	4	4	5

0 = Worst Water Quality, high incidence of algae

5 = Water Quality good, no incidence of algae.

EXAMPLE 2

Example 2 illustrates the suppression of algae in hydroponic systems.

[0025] The hydrophobic composition of Example 1 was supplemented with
10 ground cotton stubble. The composition was then applied to the drainage canals of a
seven acre hydroponics facility in Florence, Oregon, USA at a rate of 1.5 g/m². The
water quality of the hydroponics facility increased at a dramatic rate after introduction of
the hydrophobic composition. The results of the study are illustrated in Table 2.

TABLE 2

Treatment Rate	Day 0	Day 3	Day 7	Day 10
1.5g/m ²	1	2.5	4	4.5

0 = Worst Water Quality, high incidence of algae

5 = Water Quality good, no incidence of algae.

5

EXAMPLE 3

[0026] Example 3 illustrates the uniform distribution of the hydrophobic composition in a water application.

[0027] A water column was set up in Reedley, CA, USA. Various strains of *Bacillus subtilis* were formulated with charcoal to produce a hydrophobic composition with 4.5×10^8 cfu/g of hydrophobic composition.. The hydrophobic composition was then applied to the center of the surface of water as a small pile, at a range of rates from 0.1 -1.5 g/m². The rate at which the material spread across the water surface and through the column is described in Table 3. As Table 3 illustrates, the hydrophobic composition disperses rapidly over a water surface and downwards through the water column.

15

TABLE 3

Time to percentage coverage

	25%	50%	75%	100%
Water Surface	0.4 min	1.25 min	2.3 min	3.5 min
Water Column	4.75 min	7.5 min	9.5 min	12.3 min

20

EXAMPLE 4

[0028] Example 4 illustrates the effect of hydrophobic compositions on commercial turf.

[0029] The hydrophobic composition of Example 1 was applied to commercial
5 turf at Morro Bay, California, USA at rates of 2g/m^2 and 4g/m^2 . The alga incidence and the incidence of certain foliage diseases were measured 14 days later. The turf health was also measured, with 0 indicating healthy turf and 5 indicating unhealthy turf. The results of this study are illustrated in Table 4. As the rates of application of the formulation increased, the incidence of algae and leaf disease decreased, while the health
10 of the turf increased.

TABLE 4

Rates	Algae Incidence	Leaf Disease Incidence*	Turf Health**
Control	15 % increase	22 % increase	2.3
2g/m^2	-33%	- 55 %	1.5
4g/m^2	-65 %	- 75 %	1

15 * Main disease of turf at this site was leaf anthracnose caused by *Collectotrichum*.

** Turf health Index 0= no necrotic patches, growth excellent, color excellent.

5= Dead

EXAMPLE 5

20 [0030] Example 5 illustrates the use of hydrophobic compositions in aquariums and fish tanks.

[0031] Algae growth in aquariums and fish tanks are a common problem. The hydrophobic compositions of this invention may be applied to aquariums and fish tanks to treat the water and thus control the algae growth.

[0032] The hydrophobic composition of Example 1 may be placed on the water surface of an aquarium or fish tank. The hydrophobic composition rapidly disperses as a thin film over the water surface, then this film enters the liquid phase as a suspension. The *Bacillus subtilis* are being released into the body of water during dispersal over the surface of water and as the composition moves downwards through the water column to treat the algae growth in the aquarium or fish tank.

[0033] During treatment of the aquarium or fish tank for algae growth, the fish and any other aquatic/marine life forms do not need to be removed from the aquarium or fish tank. The microbial agents and hydrophobic carriers of the hydrophobic composition are not harmful to aquatic/marine life or wildlife, unlike other methods that employ chemicals or pesticides. The methods of this invention that employ microbial agents are environmentally safe.

EXAMPLE 6

[0034] Example 6 illustrates the use of hydrophobic compositions in feeding troughs for animals.

[0035] The hydrophobic composition of Example 1 may be introduced into the feeding troughs of animals. The hydrophobic composition controls the growth of target pests and diseases. Further, the hydrophobic composition is not harmful to the animals. As a result, the frequency with which the feeding troughs need to be cleaned and replaced may be decreased.

[0036] Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following claims.